

## Egg and cholesterol intake and incident type 2 diabetes among French women

Martin Lajous<sup>1,2,3</sup>, Anne Bijon<sup>3,4</sup>, Guy Fagherazzi<sup>3,4</sup>, Beverley Balkau<sup>3,4</sup>, Marie-Christine Boutron-Ruault<sup>3,4</sup> and Françoise Clavel-Chapelon<sup>3,4\*</sup>

<sup>1</sup>Center for Research on Population Health, INSP (Instituto Nacional de Salud Pública), Cuernavaca, Mexico

<sup>2</sup>Department of Epidemiology, Harvard School of Public Health, Boston, MA, USA

<sup>3</sup>Center for Research in Epidemiology and Population Health (CESP), Institut National de la Santé et de la Recherche Médicale (INSERM), Villejuif 94805, France

<sup>4</sup>Paris-South University, Orsay 91400, France

(Submitted 22 September 2014 – Final revision received 10 June 2015 – Accepted 1 July 2015 – First published online 10 September 2015)

### Abstract

Egg consumption is a major source of dietary cholesterol, a nutrient that may disrupt glucose metabolism. We prospectively evaluated the relation between egg consumption and cholesterol-intake and diabetes in 65 364 French disease-free women who responded to a validated diet history questionnaire in 1993. Egg consumption included hardboiled eggs and eggs consumed in an omelette or a mixed dish, and dietary cholesterol was estimated using a French nutrient database. Over 14 years of follow-up, 1803 incident diabetes cases were identified through self-reports, supplementary questionnaires and drug reimbursement information. Multivariable Cox regression models were adjusted for age, education, menopause, menopausal hormone therapy, hypertension and hypercholesterolaemia, BMI, physical activity, smoking, alcohol, fruit, vegetables, processed red meat, coffee and sugar and artificially sweetened beverages. No association was observed between egg consumption and risk of type 2 diabetes. When comparing women who consumed at least five eggs per week with non-consumers, the multivariable hazard ratio (HR) was found to be 1.00 (95% CI 0.78, 1.29; across categories,  $P_{\text{trend}} = 0.11$ ). Women in the highest quintile of dietary cholesterol had a 40% higher rate of diabetes compared with those in the lowest quintile (HR 1.40; 95% CI 1.19, 1.63; across quintiles,  $P_{\text{trend}} < 0.0001$ ). A 100 mg increase of dietary cholesterol per 4184 kJ (or 1000 kcal) was associated with a 14% increase in the risk of diabetes (HR 1.14; 95% CI 1.02, 1.26). In this large prospective cohort, we observed an association between dietary cholesterol and type 2 diabetes, but no association with egg consumption. In the absence of a clear underlying mechanism and potential residual confounding, these results should be interpreted with caution.

**Key words:** Prospective studies: Diet: Eggs: Cholesterol: Type 2 diabetes

The observation that high egg consumption is strongly associated with incident gestational diabetes suggests that egg consumption may disrupt glucose metabolism<sup>(1)</sup>. Eggs are a major source of cholesterol, and in animal models cholesterol intake disrupts glucose metabolism and increases circulating insulin<sup>(2)</sup>. In humans, dietary cholesterol appears to affect type 2 diabetes risk<sup>(3)</sup>, and elevated cholesterol intake prior to pregnancy is associated with an increased risk for gestational diabetes<sup>(4)</sup>. Few prospective studies have evaluated the relation between egg consumption and incidence of type 2 diabetes. Among middle-aged US adults, men and women who consumed at least one egg a day had a 58 and a 77% higher rate of diabetes compared with those who did not consume eggs<sup>(5)</sup>. This relation was not observed in a cohort of Adventists<sup>(6)</sup>, in older US adults<sup>(7)</sup>, or in two European<sup>(8,9)</sup> and a Japanese<sup>(10)</sup> population. Such inconsistency could be explained by differences in the distribution of egg consumption, in the variables used for adjustment for confounding, in the method of preparation or in

the foods consumed in conjunction with eggs. Therefore, we investigated the relation between egg and cholesterol consumption and incident type 2 diabetes in a large prospective cohort of French women.

### Methods

#### Study population

The E3N study (Etude Epidémiologique auprès des femmes de la Mutuelle Générale de l'Education Nationale) is a French prospective study of 98 995 women born between 1925 and 1950 and affiliated to a health insurance plan that covers mostly teachers and their spouses that began in 1990. E3N is the French component of the European Prospective Investigation into Cancer and Nutrition. Cohort members have returned mailed questionnaires on reproductive and lifestyle information and on newly diagnosed diseases in 1992, 1993, 1994, 1997, 2000, 2002,

**Abbreviation:** HR, hazard ratio.

\* **Corresponding author:** F. Clavel-Chapelon, email francoise.clavel@gustaveroussy.fr

2005 and 2008. Loss to follow-up has been low (3%), and average follow-up in each questionnaire cycle has been 83%. All women signed an informed consent form in compliance with the French National Commission for Computerized Data and Individual Freedom. In 1993, 74 154 women responded to a follow-up questionnaire and a validated self-administered diet history questionnaire<sup>(11)</sup>. We excluded those with an unrealistic energy consumption ( $n$  1366), no follow-up ( $n$  720), prevalent diabetes ( $n$  1576), diabetes with no date of diagnosis ( $n$  81), prevalent cancer or CVD ( $n$  4950) and women who reported following a diet for diabetes ( $n$  97). The final study population comprised 65 364 women.

### Dietary and covariate assessment

Between 1993 and 1995, dietary data were collected using a previously validated 208-item self-administered diet history questionnaire, with eleven categories of frequency: never or less than once a month; one, two or three times a month; and one to seven times a week. Total egg consumption was based on the number and frequency of consumption of hardboiled eggs and of eggs consumed in omelettes or mixed dishes. Nutrient intakes, including that of dietary cholesterol, were calculated using a food composition table derived from a French national database. The validity and reproducibility of the dietary questionnaire was previously assessed in a sample of 119 women who had completed two diet history questionnaires and twelve monthly 24-h dietary recalls over a 1-year period<sup>(11)</sup>. The correlation coefficient between the dietary questionnaire and the 24-h recalls was 0.49 for eggs and 0.40 for dietary cholesterol. Education was assessed in 1990 at cohort recruitment with the following categories: no studies, primary-school certificate, secondary-school certificate, high-school certificate (baccalaureate) to 2-year post-high-school education, 3 to 4 years of post-high-school education and 5 or more years of post-high-school education. We defined university-educated individuals as those who responded to the last two categories. For other covariates, we used information from a questionnaire sent to participants along with the dietary assessment. Treated hypertension and hypercholesterolaemia, and smoking were based on self-reports. Menopausal status was determined using information on last menstruation, the presence of hot flushes, and a history of hysterectomy, ovariectomy and menopausal hormonal treatments. Self-reported height and weight were used to calculate BMI, defined as weight (kg) divided by height squared ( $m^2$ ). In this cohort, self-reported anthropometry is reliable (in a sub-study the correlation coefficient between self-reported and measured height was 0.89  $m^2$  and for weight was 0.94 kg)<sup>(12)</sup>. We assessed usual physical activity with a questionnaire that included items on weekly hours spent walking, cycling, performing light and heavy household chores and recreational activities (e.g. swimming, tennis) during the summer and the winter. This questionnaire performed well in a previous study<sup>(13)</sup>.

### Ascertainment of type 2 diabetes

As previously described<sup>(14)</sup>, for case identification, we used self-reports, supplementary questionnaires and drug reimbursement

information. Between 1993 and 2008, 2597 self-reported incident cases were confirmed either by a supplementary questionnaire or by drug reimbursement claims' database, and an additional 843 incident cases only through this database, but validated by a supplementary questionnaire. The present analysis is based on incident cases of diabetes after applying the exclusion criteria detailed above (no date of diagnosis, unavailable dietary information or unrealistic energy consumption, no follow-up, prevalent cancer or CVD and a diet for diabetes at baseline).

### Statistical analysis

Nutrients were energy-adjusted using the residual method<sup>(15)</sup>. We categorised egg consumption as 0, <1, 1–1.9, 2–4.9 and  $\geq 5$  eggs/week, and cholesterol in quintiles, and used the lowest category as the referent. To test for trend, we used the median value for each category as a continuous variable. We evaluated cholesterol consumption using a substitution model for carbohydrates that included intakes of energy, saturated fat, unsaturated fat and protein (excluding carbohydrates) in the model. We calculated person-time from the date of completion of the dietary questionnaire to the date of diagnosis of diabetes, last follow-up available or June 2008. Hazard ratios (HR) and 95% CI were estimated from Cox regression models with age as the time scale (SAS 9.1; SAS Institute Inc.). Models were adjusted for education, menopause, menopausal hormone therapy, treated hypertension and hypercholesterolaemia, physical activity, smoking, total energy, alcohol, processed red meat, coffee, fruit, vegetables, sugar and artificially sweetened beverages and BMI. We hypothesised that BMI may modify the association between dietary variables and diabetes; therefore, we tested for statistical interaction by including a cross-product term of median egg and cholesterol consumption by BMI ( $<25$ ,  $\geq 25$  kg/ $m^2$ ) in the Cox model and by performing a log-likelihood ratio test.

### Results

Mean age was 52.7 (SD 6.6) years; the prevalence of overweight was 15.5% and that of obesity was 3.2%. Mean total egg consumption was 3.5 (SD 2.8) eggs/week, and the mean weekly hardboiled and omelette/mixed-dish egg intakes were 1.0 (SD 1.5) and 2.5 (SD 2.1), respectively. Mean cholesterol intake was 378 (SD 104) mg/d, and the main food contributors of dietary cholesterol were omelette/mixed-dish egg (17%), hardboiled egg (7%), butter (4%), liver (4%), pizza and quiche (3%), poultry (3%), creams, dairy dessert and ice creams (3%), beef (3%) and cheese (3%). The correlation coefficient for dietary cholesterol with saturated fat was 0.24 ( $P$  value  $<0.0001$ ) and  $-0.26$  ( $P$  value  $<0.0001$ ) with dietary fibre. Egg consumption and cholesterol intake were directly related to intake of alcohol, processed red meat and coffee and to BMI and hypertension. We observed an inverse relation with treated hypercholesterolaemia and education (Table 1).

We identified 1803 incident diabetes cases during the study period. After multivariable adjustment, the HR comparing women who consumed  $\geq 5$  eggs/week with non-consumers was 1.00 (95% CI 0.78, 1.29), with a  $P_{\text{trend}} = 0.11$  (Table 2). The corresponding values for hardboiled egg consumption

**Table 1.** Age-standardised risk factors for diabetes risk by total egg consumption and dietary cholesterol quintiles in a cohort of French women in 1993 (Mean values and standard deviations; percentages)

	Eggs/week					Quintiles of dietary cholesterol (mg/d)														
	0		<1		1–1.9		2–4.9		≥5		1		2		3		4		5	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Current smoker (%)	13.8		13.2		12.8		13.2		15.5		12.0		12.6		12.8		13.7		16.7	
BMI (kg/m <sup>2</sup> )																				
Mean	22.1		22.3		22.5		22.9		23.4		22.1		22.5		22.8		23.1		23.6	
SD	2.7		2.8		2.9		3.1		3.4		2.7		2.9		3.0		3.1		3.5	
Treated hypertension (%)	7.5		7.5		7.6		7.8		8.3		7.0		7.5		7.6		8.1		8.8	
Hypercholesterolaemia (%)	11.3		8.0		6.4		5.1		4.3		7.7		6.3		5.1		5.2		4.7	
Current use of MHT (%)	22.1		24.5		23.4		23.5		23.3		22.8		23.4		23.9		23.9		23.6	
Physical activity (h/week)																				
Mean	6.4		6.2		6.2		6.4		6.8		6.4		6.3		6.5		6.4		6.6	
SD	8.4		9.0		7.7		7.8		8.1		8.3		7.3		8.1		7.7		8.3	
University education (%)	40.8		40.5		38.1		35.0		32.4		38.4		37.1		36.1		34.8		33.6	
Mean dietary intake	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Total energy (kJ/d)	7909	2149	8108	1988	8453	2026	9048	2155	9814	2438	8848	2552	9073	2320	9077	2192	8969	2107	8700	2026
Processed meat (servings/d)	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.4	0.6	0.5	0.3	0.3	0.4	0.3	0.5	0.4	0.6	0.4	0.6	0.4
Fruits (servings/d)	2.6	1.8	2.5	1.6	2.5	1.6	2.5	1.6	2.6	1.8	3.1	1.9	2.7	1.7	2.5	1.6	2.3	1.5	2.1	1.5
Vegetables (servings/d)	2.3	1.3	2.2	1.2	2.2	1.2	2.3	1.2	2.4	1.3	2.4	1.3	2.3	1.2	2.2	1.2	2.3	1.2	2.3	1.3
Coffee (cups/d)	2.2	2.2	2.1	2.0	2.2	2.0	2.3	2.1	2.6	2.3	2.1	2.1	2.2	2.0	2.3	2.1	2.4	2.1	2.6	2.2
Sugar-sweetened drinks (per week)	7.6	37.8	6.2	33.8	7.0	32.2	7.6	30.0	9.2	36.3	7.3	35.5	8.2	34.7	7.9	30.2	7.9	31.1	7.1	31.3
Diet drinks (per week)	2.9	24.1	1.9	20.4	2.4	25.9	2.4	24.2	4.5	37.7	2.2	24.6	2.3	24.1	2.3	23.5	2.8	26.3	4.4	36.5
Alcohol (g/d)	9.5	13.1	10.0	13.1	10.6	12.7	11.9	13.7	13.2	15.3	8.9	11.6	10.7	12.7	11.6	13.4	12.6	14.3	14.1	16.1
Cholesterol (mg/d)	247	98	276	88	320	93	395	104	540	145	–	–	–	–	–	–	–	–	–	–

MHT, menopausal hormone therapy.

Eggs, cholesterol and incident diabetes

**Table 2.** Age-adjusted and multivariable-adjusted hazard ratios (HR) of type 2 diabetes according to the number of eggs consumed per week in the Etude Epidémiologique auprès des femmes de la Mutuelle Générale de l'Education Nationale cohort study (1993–2008) (Hazard ratios and 95 % confidence intervals)

	Eggs/week									<i>P</i> <sub>trend</sub>
	0	0-1-0.9		1-1.9		2-4.9		≥5		
		HR	95 % CI	HR	95 % CI	HR	95 % CI	HR	95 % CI	
<b>Total egg consumption</b>										
Median intake	0	0.7		1.7		3.4		6.9		
Number of cases	76	153		317		770		487		
Person-years	40 152	99 051		177 678		385 945		176 307		
Age-adjusted	Ref.	0.84	0.64, 1.10	1.01	0.79, 1.30	1.17	0.92, 1.48	1.58	1.23, 2.02	<0.0001
Multivariable*	Ref.	0.86	0.65, 1.14	1.04	0.81, 1.34	1.20	0.94, 1.52	1.56	1.22, 1.99	<0.0001
+ Diet†	Ref.	0.90	0.68, 1.19	1.06	0.82, 1.36	1.17	0.92, 1.48	1.42	1.11, 1.83	<0.0001
+ Diet† + BMI	Ref.	0.84	0.64, 1.11	0.91	0.71, 1.17	0.94	0.74, 1.20	1.00	0.78, 1.29	0.11
<b>Hardboiled eggs</b>										
Median intake	0	0.5		1.5		3.0		7.0		
Number of cases	481	644		356		252		70		
Person-years	275 240	347 858		147 855		87 954		20 226		
Age-adjusted	Ref.	1.10	0.97, 1.24	1.43	1.24, 1.64	1.67	1.43, 1.95	1.95	1.52, 2.51	<0.0001
Multivariable*	Ref.	1.08	0.96, 1.21	1.36	1.18, 1.56	1.53	1.31, 1.79	1.74	1.35, 2.25	<0.0001
+ Diet†	Ref.	1.05	0.93, 1.19	1.26	1.09, 1.45	1.37	1.17, 1.60	1.51	1.17, 1.95	<0.0001
+ Diet† + omelette/mixed-dish eggs	Ref.	1.05	0.93, 1.18	1.22	1.06, 1.41	1.31	1.12, 1.54	1.45	1.11, 1.87	<0.0001
+ Diet† + omelette/mixed-dish eggs + BMI	Ref.	1.04	0.92, 1.18	1.13	0.98, 1.30	1.13	0.96, 1.30	1.16	0.89, 1.50	0.12
<b>Omelette/mixed-dish eggs</b>										
Median intake	0	0.9		1.9		3.4		6.0		
Number of cases	143	274		500		679		207		
Person-years	69 786	169 868		259 430		298 736		81 313		
Age-adjusted	Ref.	0.81	0.66, 1.00	1.00	0.83, 1.21	1.21	1.00, 1.45	1.29	1.04, 1.60	<0.0001
Multivariable*	Ref.	0.84	0.69, 1.03	1.04	0.86, 1.25	1.24	1.03, 1.49	1.32	1.06, 1.64	<0.0001
+ Diet†	Ref.	0.87	0.71, 1.06	1.04	0.86, 1.26	1.19	0.99, 1.44	1.22	0.98, 1.52	<0.0001
+ Diet† + hardboiled eggs	Ref.	0.86	0.70, 1.06	1.02	0.85, 1.24	1.15	0.95, 1.38	1.13	0.91, 1.42	0.002
+ Diet† + hardboiled eggs + BMI	Ref.	0.85	0.69, 1.04	0.92	0.69, 1.04	0.98	0.81, 1.18	0.91	0.73, 1.14	0.67

Ref., referent values.

\* Adjusted for education, smoking (never, secondhand smoke, past, current), physical activity (h/week, quartiles), hormone replacement therapy (premenopausal, current, past, never), hypertension, hypercholesterolaemia and energy (quintiles).

† Additional adjustment alcohol (quintiles), processed red meat (quintiles), coffee (quintiles), fruits (quintiles), vegetables (quintiles), sugar-sweetened drinks (no consumers+quartiles) and artificially sweetened drinks (no consumers + quartiles).



present study the median intake for the highest quintile of intake was 507 mg/d. A recent analysis in the smaller Kuopio prospective cohort did not show an association between cholesterol intake, assessed using 4-d food records and incident diabetes after a 19-year follow-up<sup>(9)</sup>. Individuals may have changed their diet over the follow-up period, potentially attenuating the association. In a sensitivity analysis restricted to the first 10 years of follow-up, cholesterol intake appeared to confer risk of diabetes; however, results were not statistically significant. For egg consumption, our results differ from observations reported in a meta-analysis based on four prospective US studies<sup>(19)</sup> that found an association between egg consumption and type 2 diabetes and also from a study on maternal diet and gestational diabetes<sup>(1)</sup>. More recently, this association was evaluated in a small Spanish cohort study with relatively few incident cases of diabetes<sup>(8)</sup> and in the Kuopio study in Finland<sup>(9)</sup>; no association was observed in either study. Eggs are often consumed with other foods, and there are several methods of preparation that include the addition of other nutrients. Thus, other foods or added nutrients may be responsible for the observed relation. In our study, we did not observe a relation between egg consumption and diabetes risk after adjustment for BMI, a variable that is known to be strongly associated with diabetes and is also associated with diet.

The prospective design, limited loss to follow-up, use of a validated dietary questionnaire and the possibility of evaluating different egg preparations separately are important strengths of the present analysis. There are some limitations to consider. First and foremost, there is some evidence for potential biological mechanisms supporting altered cholesterol metabolism and diabetes risk. However, it is less clear how dietary cholesterol could affect diabetes occurrence. Diet was assessed only at baseline, and participants may have changed their diet during follow-up. Thus, a non-differential measurement error is possible and could potentially explain null results. Other dietary analyses in this cohort have shown significant associations with diabetes and could be indicative that this error may not be substantial. Non-differential misclassification of diabetes is also a possibility. However, assuming very few false positives, this error probably had no measurable effect on our estimates. Confounding by unmeasured factors such as dietary factors prior to baseline, or by variables that are difficult to measure, such as unhealthy behaviour, can never be ruled out. We adjusted for the most important risk factors for diabetes and saw a large attenuation of HR. We further explored this possibility by stratifying analyses on hypercholesterolaemia and hypertension, two diagnoses that could be related to unhealthy behaviours and could also alter dietary habits. However, the association between dietary cholesterol and diabetes remained among individuals without these diagnoses.

Our results suggest that dietary cholesterol is associated with type 2 diabetes, but habitual egg consumption is not. These results should be interpreted with caution as the underlying mechanism for the relation of dietary cholesterol and diabetes is unclear, and there is potential for residual confounding. Future studies should address potential mechanisms by which glucose metabolism may be disrupted by cholesterol intake.

## Acknowledgements

The authors are indebted to the participants in the Etude Epidémiologique auprès des femmes de la Mutuelle Générale de l'Education Nationale (E3N) for their continuing dedication and support. The authors thank Thierno Mamadou Bah for additional statistical support.

The E3N cohort study was carried out with the financial support of the Mutuelle Générale de l'Education Nationale, European Community, French League against Cancer, Gustave Roussy Institute and French Institute of Health and Medical Research. The validation of potential diabetes cases was supported by the European Union (Integrated Project LSHM-CT-2006-037197 in the Framework Programme 6 of the European Community) InterAct project. The present analysis was conducted with the support of the International Associated Laboratory in Nutrition, Obesity, Hormones and Cardiometabolic Disease and Depression, an ongoing collaboration between the National Institute of Health and Medical Research (Inserm, France) and the National Institute of Public Health (INSP, Mexico). M. L. is a Bernard Lown Visiting Scholar in Cardiovascular Health at the Harvard School of Public Health.

M. L. formulated the research question. M. L., A. B., M.-C. B.-R. and F. C.-C. designed the study; M. L., A. B., M.-C. B.-R. and G. F. conducted research; A. B. analysed the data; M. L., B. B. and M.-C. B.-R. and F. C.-C. wrote the paper; M. L. had primary responsibility for final content. All authors read and approved the final manuscript.

There are no conflicts of interest.

## References

1. Qiu C, Frederick IO, Zhang C, *et al.* (2011) Risk of gestational diabetes mellitus in relation to maternal egg and cholesterol intake. *Am J Epidemiol* **173**, 649–658.
2. Adamopoulos PN, Papamichael CM, Zampelas A, *et al.* (1996) Cholesterol and unsaturated fat diets influence lipid and glucose concentrations in rats. *Comp Biochem Physiol B Biochem Mol Biol* **113**, 659–663.
3. Salmeron J, Hu FB, Manson JE, *et al.* (2001) Dietary fat intake and risk of type 2 diabetes in women. *Am J Clin Nutr* **73**, 1019–1026.
4. Bowers K, Tobias DK, Yeung E, *et al.* (2012) A prospective study of prepregnancy dietary fat intake and risk of gestational diabetes. *Am J Clin Nutr* **95**, 446–453.
5. Djousse L, Gaziano JM, Buring JE, *et al.* (2009) Egg consumption and risk of type 2 diabetes in men and women. *Diabetes Care* **32**, 295–300.
6. Vang A, Singh PN, Lee JW, *et al.* (2008) Meats, processed meats, obesity, weight gain and occurrence of diabetes among adults: findings from Adventist Health Studies. *Ann Nutr Metab* **52**, 96–104.
7. Djousse L, Kamineni A, Nelson TL, *et al.* (2010) Egg consumption and risk of type 2 diabetes in older adults. *Am J Clin Nutr* **92**, 422–427.
8. Zazpe I, Beunza JJ, Bes-Rastrollo M, *et al.* (2013) Egg consumption and risk of type 2 diabetes in a Mediterranean cohort; the Sun project. *Nutr Hosp* **28**, 105–111.
9. Virtanen JK, Mursu J, Tuomainen TP, *et al.* (2015) Egg consumption and risk of incident type 2 diabetes in men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Am J Clin Nutr* **101**, 1088–1096.



10. Kurotani K, Nanri A, Goto A, *et al.* (2014) Cholesterol and egg intakes and the risk of type 2 diabetes: the Japan Public Health Center-based Prospective Study. *Br J Nutr* **112**, 1636–1643.
11. van Liere MJ, Lucas F, Clavel F, *et al.* (1997) Relative validity and reproducibility of a French dietary history questionnaire. *Int J Epidemiol* **26**, Suppl. 1, S128–S136.
12. Tehard B, van Liere MJ, Com Nougue C, *et al.* (2002) Anthropometric measurements and body silhouette of women: validity and perception. *J Am Diet Assoc* **102**, 1779–1784.
13. Tehard B, Friedenreich CM, Oppert JM, *et al.* (2006) Effect of physical activity on women at increased risk of breast cancer: results from the E3N cohort study. *Cancer Epidemiol Biomarkers Prev* **15**, 57–64.
14. Lajous M, Bijon A, Fagherazzi G, *et al.* (2014) Body mass index, diabetes, and mortality in french women: explaining away a 'paradox'. *Epidemiology* **25**, 10–14.
15. Willett W (1998) *Nutritional Epidemiology*, 2nd ed. vol. 30, Monographs in Epidemiology and Biostatistics. New York, NY: Oxford University Press.
16. Wu L, Vikramadithyan R, Yu S, *et al.* (2006) Addition of dietary fat to cholesterol in the diets of LDL receptor knockout mice: effects on plasma insulin, lipoproteins, and atherosclerosis. *J Lipid Res* **47**, 2215–2222.
17. Brunham LR, Kruit JK, Verchere CB, *et al.* (2008) Cholesterol in islet dysfunction and type 2 diabetes. *J Clin Invest* **118**, 403–408.
18. Lajous M, Tondeur L, Fagherazzi G, *et al.* (2012) Processed and unprocessed red meat consumption and incident type 2 diabetes among French women. *Diabetes Care* **35**, 128–130.
19. Shin JY, Xun P, Nakamura Y, *et al.* (2013) Egg consumption in relation to risk of cardiovascular disease and diabetes: a systematic review and meta-analysis. *Am J Clin Nutr* **98**, 146–159.